



ELSEVIER

Generation and maintenance of memory CD4⁺ T Cells

Ester MM van Leeuwen^{1,3}, Jonathan Sprent² and Charles D Surh¹

In the course of an immune response to an infectious microbe, pathogen-specific naïve CD4⁺ T cells proliferate extensively and differentiate into effector cells. Most of these cells die rapidly, but a small fraction of effector cells persist as memory cells to confer enhanced protection against the same pathogen. Recent advances indicate that strong TCR stimulation during the primary response is essential for the generation of long-lived memory CD4⁺ T cells. Memory cells appear to be derived equally from all subsets of effector cells, and memory cells can also acquire additional functional capabilities during the secondary response. Resting memory CD4⁺ cells are dependent on signals from contact with IL-7 and IL-15, but not MHC class II, for their survival and intermittent homeostatic proliferation.

Addresses

¹ The Scripps Research Institute, 10550 N. Torrey Pines Road, La Jolla, CA 92037, USA

² The Garvan Institute of Medical Research, 384 Victoria St, Darlinghurst, NSW 2010, Australia

³ Present address: Academisch Medisch Centrum, Universiteit van Amsterdam, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands.

Corresponding author: Surh, Charles D (csurh@scripps.edu)

Current Opinion in Immunology 2009, 21:167–172

This review comes from a themed issue on
Lymphocyte Development
Edited by Casey Weaver and Alexander Rudensky

Available online 11th March 2009

0952-7915/\$ – see front matter
© 2009 Elsevier Ltd. All rights reserved.

DOI 10.1016/j.coi.2009.02.005

Introduction

CD4⁺ T cells are essential players in adaptive immunity, providing help to other subsets of lymphocytes as well as mediating direct effector functions through cell-to-cell contact and via production of cytokines and chemokines [1]. Such multi-tasking capability is a reflection of the fact that functionally distinct populations of effector CD4⁺ T cells can emerge during an immune response, depending on the quality of the inflammatory conditions encountered. So far, Th1, Th2, T follicular helper (T_{FH}) and Th17 cells have been characterized as distinct subsets of effector CD4⁺ T cells [1], but whether these effector cells maintain their polarized state upon differentiation into resting memory cells is only beginning to be understood. Although studies on CD4⁺ T cell responses have been facilitated by the use of TCR

transgenic (Tg) cells and MHC class II (MHC-II) tetramers, memory CD4⁺ T cells are still less well understood than their CD8⁺ counterparts. While the two subsets of T cells share some common characteristics, the mechanisms involved in the formation of CD4⁺ memory T cells are different from CD8⁺ memory T cells [2]. Here, we review recent advances in defining the mechanisms involved in the generation and maintenance of memory CD4⁺ T cells.

TCR signalling intensity and memory CD4⁺ T cell development

Unlike naïve CD8⁺ T cells that can be easily triggered into a programmed course of differentiation with a relatively brief period of stimulation [3,4], naïve CD4⁺ T cells require prolonged antigenic stimulation in order to differentiate into effector and memory cells [5,6]. Moreover, while past studies have suggested that weaker stimulation is better than strong stimulation for memory formation [7,8], more recent studies indicate that naïve CD4⁺ T cells require a high threshold of TCR signalling in order to fully differentiate into effector cells that can convert to memory cells. Much of the evidence supporting this notion emerged from the finding that the presence of an unphysiologically high precursor frequency of antigen (Ag)-specific naïve CD4⁺ T cells during the primary response is detrimental for memory formation. Similar findings were reported previously for CD8⁺ T cells [9,10], but it is now apparent that the influence of precursor frequency on memory cell generation is much greater for CD4⁺ cells than CD8⁺ cells. The negative effect of high precursor frequency on the quality of CD4⁺ memory cells was initially reported for ovalbumin (ova)-specific OT-II Tg cells and shown to impair their ability to undergo proliferation and secrete cytokines in response to *Listeria monocytogenes* (Lm) expressing ova [11]. A more thorough study with E α -specific TE α Tg cells revealed not only that the response to vesicular stomatitis virus (VSV) expressing E α at a high precursor frequency led to the generation of effector cells with impaired ability to secrete cytokines but also that these cells were unable to survive as memory cells [12[•]]. Moreover, this defective immune response was mimicked when a low precursor frequency of naïve TE α Tg cells (which yields a normal immune response) was exposed to only limited amounts of Ag, that is, by injecting a mAb that competes for the TCR ligand [12[•]].

Further evidence for the requirement for a high threshold of TCR stimulation for memory generation emerged from studies with Smarta1 TCR Tg CD4⁺ T cells specific for

lymphocytic choriomeningitis virus (LCMV) glycoprotein (gp) 61 presented by LCMV versus Lm [13[•]]. Unexpectedly, although naïve Smarta1 cells underwent a comparable level of expansion to both LCMV and Lm expressing gp61 (Lm-gp61), memory CD4⁺ T cells developed only with LCMV and not with Lm-gp61 infection. The failure to generate memory cells with Lm-gp61 occurred regardless of the T cell precursor frequency, indicating that clonal competition for Ag was not the issue [13[•]]. Rather, it appeared that Lm-gp61 infection led to a subthreshold expression of Ag density, as suggested by the findings that Lm-gp61-induced effector cells displayed reduced effector function and a lower TCR avidity than control LCMV-induced effector cells [13[•]]. Even when the density of the Ag was not limiting, as in the case with LCMV infection, high precursor frequencies of Smarta1 cells led to the generation of memory CD4⁺ T cells with an abbreviated lifespan and impaired ability to mount a secondary response [14[•]]. Similar shortening in the lifespan of memory CD4⁺ cells was also observed for two other lines of TCR Tg CD4⁺ cells when they were stimulated with peptide and adjuvant at a high precursor frequency [15]. Interestingly, in the case with Smarta1 cells, competition for cytokines, in particular IFN γ [14[•]], appeared to be the main cause of aberrant memory cell development. Collectively, these reports indicate that strong TCR signalling, together with appropriate costimulatory signals, is required for optimal generation of long-lived memory CD4⁺ T cells.

Recent intravital two-photon imaging studies provided a cellular mechanism for why clonal competition at a high precursor frequency of Ag-specific CD4⁺ T cells is detrimental for a normal immune response. Here, an elegant system was devised to observe the initial interaction of T cells with Ag-loaded dendritic cells (DC). The approach involved reconstituting MHC-II⁻ mice with MHC-II⁺ DC and TCR Tg CD4⁺ T cells expressing different markers and then imaging the draining LN right after intravenous injection of the agonist peptide [16[•]]. In contrast to past reports of only transient interactions between T cells and DC during the first few hours of Ag recognition [17], prolonged T-DC interactions were observed immediately in this system and lasted for several hours [16[•]]. By terminating the T-DC interaction at various time points with an injection of anti-MHC-II mAb, it was found that at least six hours of continuous contact was required for naïve CD4⁺ T cells to enter the cell cycle and more than 24 h for further cell proliferation and synthesis of cytokines [16[•]]. Importantly, prolonged T-DC interactions were readily observed with a low precursor frequency of Ag-specific CD4⁺ T cells for the first 48 h, whereas the duration of T-DC interaction became much shorter, as previously reported, at a high precursor frequency of naïve CD4⁺ T cells [18[•]]. Thus, clonal competition at a high precursor frequency appears to impair the ability of T cells to make prolonged meaningful contacts with Ag-loaded DC.

Linear versus non-linear differentiation of memory CD4⁺ T cells

There are two competing models of memory T cell development: linear and divergent differentiation. In the first model, activated naïve T cells develop initially into cytokine-producing effector cells before a small fraction of these cells convert to memory cells. In the divergent model, a fraction of activated naïve T cells differentiate directly into memory cells, bypassing the effector phase. A report supporting the divergent model appeared several years ago with the finding that purified IFN γ ⁺ TCR Tg effector CD4⁺ T cells generated with peptide immunization failed to survive upon adoptive transfer into naïve hosts, whereas IFN γ ⁻ Tg cells persisted as memory cells and produced IFN γ upon re-stimulation [19].

Two recent studies on the response to LCMV infection, however, are more consistent with the linear differentiation model. Using a similar approach to the previous report, one study found that generation of memory cells was equally efficient for IFN γ ⁺ and IFN γ ⁻ Smarta1 effector cells upon adoptive transfer into naïve hosts [20^{••}]. The other study used two complementary genetic approaches to mark IFN γ ⁺ polyclonal effector T cells with CD90.1 (Thy-1.1) and then studied the fate of CD90.1⁺ and CD90.1⁻ cells upon adoptive transfer into naïve hosts. Again, IFN γ ⁺ effector CD4⁺ (and CD8⁺) T cells were found to differentiate into memory cells just as efficiently, or even more efficiently, than IFN γ ⁻ effector cells [21^{••}]. Both reports found that memory cells derived from IFN γ ⁺ effectors were fully functional by all parameters analyzed and also confirmed that memory cells generated from IFN γ ⁻ effectors produced IFN γ upon re-stimulation. Therefore, these reports provide strong evidence that development of memory CD4⁺ T cells, as with memory CD8⁺ T cells, occurs through a linear pathway. Since IFN γ ⁻ effector cells gave rise to memory cells, it is also possible that the divergent pathway may contribute to memory CD4⁺ T cell formation. One possible indication of the divergent pathway is that recently activated naïve CD8⁺ T cells initially undergo asymmetric cell division, the proximal daughter cell displaying both functional and phenotypic characteristics of effector cells and the distal daughter cell behaving more like a memory cell [22]. Asymmetric cell division was also observed for CD4⁺ T cells, although whether distal daughter cells also exhibit memory function was not analyzed.

IL-7 and selection of memory CD4⁺ T cells

Signalling through the IL-7 receptor (IL-7R), a heterodimer of IL-7R α and the common γ chain (γ_c), is essential for prolonged survival of naïve and memory T cells. IL-7R is normally expressed at high levels on resting T cells, and IL-7R α is rapidly downregulated upon T cell activation and re-expressed by a small fraction of effector T

cells destined to survive as memory cells [23]. Recent work suggests that contact with IL-2 during the early phase of activation promotes IL-7R α re-expression on CD4⁺ effector cells [24]. Nonetheless, for CD8⁺ T cells it is now clear that IL-7R expression, while useful for detecting memory precursors, is not required for the effector-to-memory cell transition [25–27]. For CD4⁺ T cells, however, IL-7R signalling has a much bigger role, and appears to be essential, though not sufficient, for development of memory CD4⁺ cells. The requirement for contact with IL-7 is suggested by the finding that memory CD4⁺ cells fail to arise when immune responses are generated in IL-7⁻ hosts and with T cells expressing a mutant form of IL-7R α (IL-7R α ^{449F}) that cannot induce STAT5 activation [28,29]. Others examined the role of IL-7R signalling at different phases of the CD4⁺ T cell response. One study used polyclonal CD4⁺ T cells expressing chimeric GM-CSF/IL-7R α chain to assess the effect of strong IL-7R signalling during the early phase of the immune response to LCMV, thereby eliciting intense IL-7R signalling delivered via the burst of GM-CSF produced during the first six days after infection. While this strategy increased the magnitude of the effector cell response, the contraction phase was reciprocally pronounced, thus resulting in no enhancement in the overall production of memory cells [26]. Nonetheless, another study found that provision of strong IL-7 signalling by administration of long-lived IL-7/mAb complexes at the peak of the response prevented the contraction of TCR Tg effector CD4⁺ cells by increasing their homeostatic proliferation and Bcl-2 upregulation [30^{*}]. Although these findings indicate that signalling through IL-7 can augment the immune response, constitutive expression of Tg IL-7R α on T cells failed to enhance (or impair) the production of effector and memory CD4⁺ (and CD8⁺) cells [27,31^{*}] and their ability to display normal secondary responses [31^{*}]. Thus, continued expression of IL-7R α alone is not sufficient for the development of memory CD4⁺ T cells.

Homeostasis of memory CD4⁺ T cells

While most investigators agree that signalling from a combination of IL-7 and IL-15, but not MHC-I, regulates homeostasis of memory CD8⁺ T cells, the factors controlling homeostasis of the memory CD4⁺ T cell pool are more controversial [32]. Much of the confusion here seems to stem from the invalid assumption that memory-phenotype (MP) CD4⁺ T cells, which arise spontaneously without intentional immunization, and Ag-specific (AgSp) memory CD4⁺ T cells, which arise after intentional immunization with a specific Ag, display the same homeostatic requirements (see below). Recent studies with AgSp TCR Tg and polyclonal memory CD4⁺ cells demonstrate that these cells, like memory CD8⁺ T cells, rely on contact with both IL-15 and IL-7 but not MHC (MHC-II) for their homeostasis [28,33^{**},34^{*}]. IL-7 appears to have a bigger role than IL-15 in supporting

survival of memory CD4⁺ cells, but both cytokines seem equally essential for these cells to undergo basal homeostatic proliferation under normal conditions [33^{**},34^{*}]. IL-15 has a less prominent role for memory CD4⁺ cells than for NK and memory CD8⁺ cells, these latter cells being much more reliant on IL-15 than IL-7 for their homeostasis [32]. This difference in IL-15 dependence correlates closely with the expression levels of the IL-15 receptor, CD122, which is displayed at much lower levels on memory CD4⁺ cells than on NK and memory CD8⁺ cells. For this reason, memory CD4⁺ cells compete less effectively for IL-15 than NK and memory CD8⁺ cells [33^{**}], and this could be one of the main reasons why the lifespan of AgSp memory CD4⁺ cells tends to be shorter than that of memory CD8⁺ cells [35].

In contrast to AgSp memory CD4⁺ cells, MP CD4⁺ cells comprise a heterogeneous population of cells with different homeostatic requirements. While the majority of MP cells appear to resemble AgSp memory CD4⁺ cells, a small fraction of MP CD4⁺ cells undergoes a very rapid rate of homeostatic proliferation under both normal and lymphopenic conditions. Significantly, such rapid proliferation is driven largely by TCR interaction with MHC-II and not by contact with IL-7 or IL-15 [33^{**},36]. The presence of a population of fast-dividing MP CD4⁺ cells has been interpreted to indicate a role for TCR-MHC-II signalling in homeostasis of memory CD4⁺ cells [37]. However, this conclusion is questionable because AgSp memory CD4⁺ cells typically divide quite slowly [28,33^{**},34^{*}]. The function of the fast-dividing subset of MP CD4⁺ cells and the nature of the Ag that drives their rapid proliferation have yet to be defined.

Role of apoptotic proteins in memory T cell survival

A delicate balance of pro-apoptotic and anti-apoptotic molecules dictates survival of T cells [38]. Both IL-7 and IL-15 have been shown to support T cell survival by increasing the expression of anti-apoptotic molecules, such as Bcl-2 and Mcl-1 [39]. On the contrary, the pro-apoptotic BH3-only molecule Bim has been implicated in effector cell death during the contraction phase [40]. This notion was recently confirmed for CD4⁺ T cells with the finding that a larger pool of effector cells survived to differentiate into memory cells in Bim^{-/-} mice than in normal mice; however, Bim⁻ memory CD4⁺ cells did not have an overtly extended lifespan, implying that other homeostatic mechanisms beside Bim are involved in mediating death of memory cells [41]. In addition, high levels of Bim expression were found on effector CD4⁺ T cells derived from suboptimally activated naïve CD4⁺ T cells, that is, cells that failed to develop into memory cells [13^{*}].

The role of Bcl-2 in CD4⁺ memory cell generation has been studied in Bim^{+/-} Bcl-2^{-/-} mice, thus circumventing the

problem of early death of Bcl-2^{-/-} mice [42[•]]. In contrast to the severe loss of naïve T cells in Bim^{+/-} Bcl-2^{-/-} mice, production of LCMV-specific memory CD4⁺ cells was only slightly reduced. However, much of the memory CD4⁺ cell production in these mice appeared to reflect enhanced lymphopenia-driven proliferation; thus, these memory CD4⁺ cells rapidly disappeared upon adoptive transfer into normal T-cell-sufficient hosts [42[•]]. The conclusion therefore is that Bcl-2 has a prominent role in controlling the survival of memory CD4⁺ T cells.

In a study using human T cells, gene expression profiling and proteomic analysis suggested that survival of human memory CD4⁺ T cells is mediated partly by inhibiting the transcription of Bim and Fas via inactivation of forkhead box O3a (FOXO3a) transcription factor combined with activation of STAT5a [43[•]]. Indeed, silencing the active form of FOXO3a by introducing small interfering RNA or a dominant negative form of FOXO3a was recently found to enhance survival of memory CD4⁺ T cells in HIV patients [44[•]]. Another BH3-only pro-apoptotic molecule, Noxa, was also reported to play an intermediary role in controlling memory CD4⁺ T cell generation [45]. Although the absence of Noxa itself did not affect development of naïve and memory T cells, significant defects were seen with loss of the polycomb group gene, Bmi1, which controls Noxa expression. Thus, CD4⁺ T cells deficient in Bmi1 underwent normal proliferation and differentiation into effector cells but were severely impaired in their ability to further differentiate into memory cells [46]. Interestingly, the premature death of Bmi1⁻ effector cells appeared to be mediated through increased Noxa expression, indicating that Bmi1 promotes memory CD4⁺ cell generation through repression of the Noxa gene.

Recall response of memory CD4⁺ T cells

Rapid recall immune responses by memory T cells are attributed largely to their increased precursor frequency of Ag-specific cells plus their heightened responsiveness to Ag. One of the major reasons for increased sensitivity to Ag is the reduced co-stimulatory requirements for activation of memory T cells compared to naïve T cells. The poised state of memory CD4⁺ T cells also appears to reflect elevated expression of the TCR proximal tyrosine kinase, Zap70, in memory CD4⁺ T cells [47]. With regard to co-stimulatory signals, recent work confirmed that these signals are not required for the initial activation of memory CD4⁺ T cells to Ag, but nonetheless are important for activated memory cells to mount a sustained immune response. Thus, *in vivo* blocking of CD28/B7 interactions by injection of CTLA4-Ig considerably reduced the magnitude of memory cell expansion and impaired the capacity of the expanded cells to secrete IL-2 and IFN γ [48]. Moreover, another group found that while CD40/CD40L interactions were dispensable for the expansion of memory CD4⁺ T cells, CD40L signals

were nevertheless required for these cells to synthesize IFN γ [49].

With regard to the kinetics of memory cell proliferation, recent work on viral infections has confirmed the long-held belief that naïve and memory CD4⁺ T cells commence cell proliferation at a similar time point after Ag exposure and undergo a similar rate of cell division [50]. Thus, LCMV-specific Smarta1 memory CD4⁺ T cells, like naïve Smarta1 cells, did not initiate cell division until three to four days after LCMV infection, despite the fact that memory cells synthesized IFN γ within several hours after the infection. The reason for the lag before entry into cell division is not clear but seems to be dictated by the microenvironment of the infected host. Thus, transfer of Smarta1 cells into mice pre-infected with LCMV two days before entered cell division earlier [50]. In light of these findings, the higher magnitude of the secondary response compared to the primary response can be attributed largely to the increased precursor frequency of Ag-specific cells. Nonetheless, another study found that LCMV-specific polyclonal memory CD4⁺ cells proliferated at a slower rate than naïve CD4⁺ cells towards the end of the response [51]. The reduced proliferation of memory cells appeared to be due to their low production of IL-2 and increased synthesis of IFN γ , as provision of exogenous IL-2 or anti-IFN γ mAb increased the proliferation of the memory cells [51].

Differentiation of naïve CD4⁺ T cells into Th1 or Th2 subsets of effector cells is known to involve heritable epigenetic changes that allow for expression of a restricted set of cytokines. For this reason, memory CD4⁺ cells are widely believed to maintain the polarized cytokine profile of their predecessors. This notion, however, has recently been challenged by two studies demonstrating that Th1 and Th2 memory cells can be induced to synthesize the cytokines of the alternate lineage when stimulated under the opposing conditions. One group analyzed the memory cells generated from *in vitro*-polarized Th2 (and Th1) Smarta1 effector cells that were adoptively transferred into normal hosts. While Th2 Smarta1 memory cells synthesized Th2 cytokines when left unstimulated, infection with LCMV, a strong inducer of Th1 cell development, caused the bulk of Th2 memory cells to produce IFN γ in addition to Th2 cytokines [20^{••}]. The other group used DC pulsed with an agonist peptide plus either a Th1-inducing bacterial extract or Th2-inducing worm egg Ag to generate Th1 or Th2 responses in normal mice. When the primed mice were boosted with Ag in Th1-inducing or Th2-inducing conditions, a large fraction of memory cells in both situations produced both Th1 and Th2 cytokines [52[•]]. These findings indicate that memory CD4⁺ cells derived from polarized effectors are much more amenable than previously thought and are reminiscent of a past study that found that polarized

human Th1 and Th2 cells can synthesize cytokines of the alternate lineage [53].

Concluding remarks

Much progress has been made in recent years in understanding the generation and maintenance of memory CD4⁺ T cells. It is now clear that the strength of TCR signalling during priming has a strong impact on memory formation and that memory cells can develop from both cytokine-producing and non-producing effector cells. Memory CD4⁺ cells are maintained by signals from contact with both IL-7 and IL-15, but not from TCR interaction with MHC-II ligands. Memory cells generated from polarized Th1 and Th2 effector cells are more flexible than previously considered in terms of their cytokine production capability. This finding implies that pathogenic Th cells in chronic and autoimmune disease could be manipulated to express more benign traits of a different Th lineage. Despite these advances, many key issues still remain outstanding. Two areas are particularly noteworthy. First is the age-old question of the distinguishing feature of the small fraction of effector cells that is selected to persist as memory cells. Even with recent advances, this issue remains largely an enigma. The second important question is why the lifespan of memory CD4⁺ cells is shorter than for memory CD8⁺ cells. This question is also unresolved and highlights the increasing realization that many of mechanisms involved in the generation and maintenance of memory CD4⁺ and CD8⁺ cells are subtly, but distinctly different.

Acknowledgements

This work was supported by US Public Health Service grants AI064586 and AI045809 to CDS and Australian NHMRC grants to JS. EMMvL was partly supported by a fellowship from the Netherlands Organisation for Scientific Research (NWO) and a pilot project award from The Cooperative Study Group for Autoimmune Disease Prevention (NIH). This is manuscript # 19939 from TSRI.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Swain SL, Agrewala JN, Brown DM, Jelley-Gibbs DM, Golech S, Huston G, Jones SC, Kamperschroer C, Lee WH, McKinstry KK *et al.*: **CD4⁺ T-cell memory: generation and multi-faceted roles for CD4⁺ T cells in protective immunity to influenza.** *Immunol Rev* 2006, **211**:8-22.
 2. Seder RA, Ahmed R: **Similarities and differences in CD4⁺ and CD8⁺ effector and memory T cell generation.** *Nat Immunol* 2003, **4**:835-842.
 3. Kaech SM, Ahmed R: **Memory CD8⁺ T cell differentiation: initial antigen encounter triggers a developmental program in naive cells.** *Nat Immunol* 2001, **2**:415-422.
 4. van Stipdonk MJ, Lemmens EE, Schoenberger SP: **Naive CTLs require a single brief period of antigenic stimulation for clonal expansion and differentiation.** *Nat Immunol* 2001, **2**:423-429.
 5. Williams MA, Bevan MJ: **Shortening the infectious period does not alter expansion of CD8 T cells but diminishes their capacity to differentiate into memory cells.** *J Immunol* 2004, **173**:6694-6702.
 6. Obst R, van Santen HM, Mathis D, Benoist C: **Antigen persistence is required throughout the expansion phase of a CD4⁺ T cell response.** *J Exp Med* 2005, **201**:1555-1565.
 7. Catron DM, Rusch LK, Hataye J, Itano AA, Jenkins MK: **CD4⁺ T cells that enter the draining lymph nodes after antigen injection participate in the primary response and become central-memory cells.** *J Exp Med* 2006, **203**:1045-1054.
 8. Jelley-Gibbs DM, Brown DM, Dibble JP, Haynes L, Eaton SM, Swain SL: **Unexpected prolonged presentation of influenza antigens promotes CD4 T cell memory generation.** *J Exp Med* 2005, **202**:697-706.
 9. Badovinac VP, Haring JS, Harty JT: **Initial T cell receptor transgenic cell precursor frequency dictates critical aspects of the CD8⁺ T cell response to infection.** *Immunity* 2007, **26**:827-841.
 10. Marzo AL, Klonowski KD, Le Bon A, Borrow P, Tough DF, Lefrancois L: **Initial T cell frequency dictates memory CD8⁺ T cell lineage commitment.** *Nat Immunol* 2005, **6**:793-799.
 11. Foulds KE, Shen H: **Clonal competition inhibits the proliferation and differentiation of adoptively transferred TCR transgenic CD4 T cells in response to infection.** *J Immunol* 2006, **176**:3037-3043.
 12. Blair DA, Lefrancois L: **Increased competition for antigen during priming negatively impacts the generation of memory CD4 T cells.** *Proc Natl Acad Sci U S A* 2007, **104**:15045-15050.
 - References [12^{*}-14^{*}] show that a threshold of strong TCR plus co-stimulatory signalling is required during the primary response for optimal development of effector and memory CD4⁺ T cells. The suboptimal signalling can lead to defects in both effector and memory cells or mainly to memory cells.
 13. Williams MA, Ravkov EV, Bevan MJ: **Rapid culling of the CD4⁺ T cell repertoire in the transition from effector to memory.** *Immunity* 2008, **28**:533-545.
 - Same as reference [12^{*}].
 14. Whitmire JK, Benning N, Eam B, Whitton JL: **Increasing the CD4⁺ T cell precursor frequency leads to competition for IFN-gamma thereby degrading memory cell quantity and quality.** *J Immunol* 2008, **180**:6777-6785.
 - Same as reference [12^{*}].
 15. Hataye J, Moon JJ, Khoruts A, Reilly C, Jenkins MK: **Naive and memory CD4⁺ T cell survival controlled by clonal abundance.** *Science* 2006, **312**:114-116.
 16. Celli S, Lemaitre F, Bousso P: **Real-time manipulation of T cell-dendritic cell interactions *in vivo* reveals the importance of prolonged contacts for CD4⁺ T cell activation.** *Immunity* 2007, **27**:625-634.
 - Together with reference [18^{*}], this paper demonstrates that naive CD4⁺ T cells engage continuously with DC immediately upon recognition of Ag for several hours before entering the cell cycle and that this prolonged interaction becomes rare with severe clonal competition for Ag.
 17. Mempel TR, Henrickson SE, Von Andrian UH: **T-cell priming by dendritic cells in lymph nodes occurs in three distinct phases.** *Nature* 2004, **427**:154-159.
 18. Garcia Z, Pradelli E, Celli S, Beuneu H, Simon A, Bousso P: **Competition for antigen determines the stability of T cell-dendritic cell interactions during clonal expansion.** *Proc Natl Acad Sci U S A* 2007, **104**:4553-4558.
 - Same as reference [16^{*}].
 19. Wu CY, Kirman JR, Rotte MJ, Davey DF, Perfetto SP, Rhee EG, Freidag BL, Hill BJ, Douek DC, Seder RA: **Distinct lineages of T(H)1 cells have differential capacities for memory cell generation *in vivo*.** *Nat Immunol* 2002, **3**:852-858.
 20. Lohning M, Hegazy AN, Pinschewer DD, Busse D, Lang KS, Hofer T, Radbruch A, Zinkernagel RM, Hengartner H: **Long-lived virus-reactive memory T cells generated from purified cytokine-secreting T helper type 1 and type 2 effectors.** *J Exp Med* 2008, **205**:53-61.
 - This reference along with reference [21^{**}] provides strong evidence that naive CD4⁺ T cells develop into memory cells via linear differentiation

pathway by passing through the effector cell stage. The papers clearly show that memory cells emerge equally efficiently from both cytokine-producing and non-producing effector cells.

21. Harrington LE, Janowski KM, Oliver JR, Zajac AJ, Weaver CT:
 •• **Memory CD4 T cells emerge from effector T-cell progenitors.** *Nature* 2008, **452**:356-360.
 Same as reference [20*].
22. Chang JT, Palanivel VR, Kinjyo I, Schambach F, Intlekofer AM, Banerjee A, Longworth SA, Vinup KE, Mrass P, Oliaro J *et al.*:
Asymmetric T lymphocyte division in the initiation of adaptive immune responses. *Science* 2007, **315**:1687-1691.
23. Kaech SM, Tan JT, Wherry EJ, Konieczny BT, Surh CD, Ahmed R:
Selective expression of the interleukin 7 receptor identifies effector CD8 T cells that give rise to long-lived memory cells. *Nat Immunol* 2003, **4**:1191-1198.
24. Dooms H, Wolslegel K, Lin P, Abbas AK: **Interleukin-2 enhances CD4+ T cell memory by promoting the generation of IL-7R alpha-expressing cells.** *J Exp Med* 2007, **204**:547-557.
25. Klonowski KD, Williams KJ, Marzo AL, Lefrancois L: **Cutting edge: IL-7-independent regulation of IL-7 receptor alpha expression and memory CD8 T cell development.** *J Immunol* 2006, **177**:4247-4251.
26. Sun JC, Lehar SM, Bevan MJ: **Augmented IL-7 signaling during viral infection drives greater expansion of effector T cells but does not enhance memory.** *J Immunol* 2006, **177**:4458-4463.
27. Hand TW, Morre M, Kaech SM: **Expression of IL-7 receptor alpha is necessary but not sufficient for the formation of memory CD8 T cells during viral infection.** *Proc Natl Acad Sci U S A* 2007, **104**:11730-11735.
28. Kondrack RM, Harbertson J, Tan JT, McBreen ME, Surh CD, Bradley LM: **Interleukin 7 regulates the survival and generation of memory CD4 cells.** *J Exp Med* 2003, **198**:1797-1806.
29. Osborne LC, Dhanji S, Snow JW, Priatel JJ, Ma MC, Miners MJ, Teh HS, Goldsmith MA, Abraham N: **Impaired CD8 T cell memory and CD4 T cell primary responses in IL-7R alpha mutant mice.** *J Exp Med* 2007, **204**:619-631 Epub 2007 Feb 2026.
30. Tripathi P, Mitchell TC, Finkelman F, Hildeman DA: **Cutting Edge: Limiting amounts of IL-7 do not control contraction of CD4+ T cell responses.** *J Immunol* 2007, **178**:4027-4031.
 Together with reference [31*], this paper suggests that continuous signaling through the IL-7 receptor on effector CD4+ T cells alone is not sufficient for further differentiation into memory cells.
31. Haring JS, Jing X, Bollenbacher-Reilley J, Xue HH, Leonard WJ, Harty JT: **Constitutive expression of IL-7 receptor alpha does not support increased expansion or prevent contraction of antigen-specific CD4 or CD8 T cells following *Listeria monocytogenes* infection.** *J Immunol* 2008, **180**:2855-2862.
 Same as reference [30*].
32. Surh CD, Sprent J: **Homeostasis of naive and memory T cells.** *Immunity* 2008, **29**:848-862.
33. Purton JF, Tan JT, Rubinstein MP, Kim DM, Sprent J, Surh CD:
 •• **Antiviral CD4+ memory T cells are IL-15 dependent.** *J Exp Med* 2007, **204**:951-961.
 References [33**,34*] demonstrate that homeostasis of Ag-specific memory CD4+ T cells is regulated by signals from contact with both IL-7 and IL-15, but not from MHC class II molecules. Reference [34*] also illustrates that memory-phenotype (MP) CD4+ T cells are composed of heterogeneous subsets of cells with distinct homeostatic requirements.
34. Lenz DC, Kurz SK, Lemmens E, Schoenberger SP, Sprent J,
 • Oldstone MB, Homann D: **IL-7 regulates basal homeostatic proliferation of antiviral CD4+T cell memory.** *Proc Natl Acad Sci U S A* 2004, **101**:9357-9362.
 Same as reference [33**].
35. Homann D, Teyton L, Oldstone MB: **Differential regulation of antiviral T-cell immunity results in stable CD8+ but declining CD4+ T-cell memory.** *Nat Med* 2001, **7**:913-919.
36. Robertson JM, MacLeod M, Marsden VS, Kappler JW, Marrack P:
Not all CD4+ memory T cells are long lived. *Immunol Rev* 2006, **211**:49-57.
37. Seddon B, Tomlinson P, Zamoyska R: **Interleukin 7 and T cell receptor signals regulate homeostasis of CD4 memory cells.** *Nat Immunol* 2003, **4**:680-686.
38. Marsden VS, Strasser A: **Control of apoptosis in the immune system: Bcl-2, BH3-only proteins and more.** *Annu Rev Immunol* 2003, **21**:71-105.
39. Khaled AR, Durum SK: **Lymphocide: cytokines and the control of lymphoid homeostasis.** *Nat Rev Immunol* 2002, **2**:817-830.
40. Strasser A, Pellegrini M: **T-lymphocyte death during shutdown of an immune response.** *Trends Immunol* 2004, **25**:610-615.
41. Wojciechowski S, Jordan MB, Zhu Y, White J, Zajac AJ, Hildeman DA: **Bim mediates apoptosis of CD127(lo) effector T cells and limits T cell memory.** *Eur J Immunol* 2006, **36**:1694-1706.
42. Wojciechowski S, Tripathi P, Bourdeau T, Acero L, Grimes HL,
 • Katz JD, Finkelman FD, Hildeman DA: **Bim/Bcl-2 balance is critical for maintaining naive and memory T cell homeostasis.** *J Exp Med* 2007, **204**:1665-1675.
 Together with reference [41], this paper shows that the pro-apoptotic BH3-only molecule Bim mediates death of effector, but not memory CD4+ T cells, and that Bcl-2 is not only involved in survival of naive T cells but is also essential for survival of memory CD4+ T cells.
43. Riou C, Yassine-Diab B, Van grevenynghe J, Somogyi R,
 • Greller LD, Gagnon D, Gimmig S, Wilkinson P, Shi Y, Camerer MJ *et al.*: **Convergence of TCR and cytokine signaling leads to FOXO3a phosphorylation and drives the survival of CD4+ central memory T cells.** *J Exp Med* 2007, **204**:79-91.
 References [43*,44*] identify forkhead box O3a (FOXO3a) as a transcription factor that activates pro-apoptotic mediators responsible for death of memory CD4+ T cells.
44. van Grevenynghe J, Procopio FA, He Z, Chomont N, Riou C,
 • Zhang Y, Gimmig S, Boucher G, Wilkinson P, Shi Y *et al.*: **Transcription factor FOXO3a controls the persistence of memory CD4(+) T cells during HIV infection.** *Nat Med* 2008, **14**:266-274.
 Same as reference [43*].
45. Villunger A, Michalak EM, Coultas L, Mullauer F, Bock G, Auserlechner MJ, Adams JM, Strasser A: **p53- and drug-induced apoptotic responses mediated by BH3-only proteins puma and noxa.** *Science* 2003, **302**:1036-1038.
46. Yamashita M, Kuwahara M, Suzuki A, Hirahara K, Shinnaksu R, Hosokawa H, Hasegawa A, Motohashi S, Iwama A, Nakayama T: **Bmi1 regulates memory CD4 T cell survival via repression of the Noxa gene.** *J Exp Med* 2008, **205**:1109-1120.
47. Chandok MR, Okoye FI, Ndejemi MP, Farber DL: **A biochemical signature for rapid recall of memory CD4 T cells.** *J Immunol* 2007, **179**:3689-3698.
48. Ndejemi MP, Teijaro JR, Patke DS, Bingaman AW, Chandok MR, Azimzadeh A, Nadler SG, Farber DL: **Control of memory CD4 T cell recall by the CD28/B7 costimulatory pathway.** *J Immunol* 2006, **177**:7698-7706.
49. MacLeod M, Kwakkenbos MJ, Crawford A, Brown S, Stockinger B, Schepers K, Schumacher T, Gray D: **CD4 memory T cells survive and proliferate but fail to differentiate in the absence of CD40.** *J Exp Med* 2006, **203**:897-906.
50. Whitmire JK, Eam B, Whitton JL: **Tentative T cells: memory cells are quick to respond, but slow to divide.** *PLoS Pathog* 2008, **4**:e1000041.
51. MacLeod MK, McKee A, Crawford F, White J, Kappler J, Marrack P: **CD4 memory T cells divide poorly in response to antigen because of their cytokine profile.** *Proc Natl Acad Sci U S A* 2008, **105**:14521-14526.
52. Krawczyk CM, Shen H, Pearce EJ: **Functional plasticity in memory T helper cell responses.** *J Immunol* 2007, **178**:4080-4088.
 Together with reference [20**], this paper shows that memory CD4+ T cells derived from polarized Th1 and Th2 effector cells can produce cytokines of the alternate lineage when stimulated under opposing conditions.
53. Messi M, Giacchetto I, Nagata K, Lanzavecchia A, Natoli G, Sallusto F: **Memory and flexibility of cytokine gene expression as separable properties of human T(H)1 and T(H)2 lymphocytes.** *Nat Immunol* 2003, **4**:78-86.