

# Prevalence of Osteoporosis in Thai Men

Chatlert Pongchaiyakul MD\*,  
Chalermchai Apinyanurag MD\*\*, Supasil Soontrapa MD\*\*\*,  
Sugree Soontrapa MD\*\*\*\*, Choowong Pongchaiyakul BSc\*\*\*\*\*,  
Tuan V Nguyen PhD\*\*\*\*\*, Rajata Rajatanavin MD\*\*\*\*\*

\* Departments of Medicine, Faculty of Medicine, Khon Kaen University, Khon Kaen

\*\* Departments of Radiology, Faculty of Medicine, Khon Kaen University, Khon Kaen

\*\*\* Departments of Orthopaedics, Faculty of Medicine, Khon Kaen University, Khon Kaen

\*\*\*\* Departments of Obstetric and Gynecology, Hospital, Faculty of Medicine, Khon Kaen University, Khon Kaen

\*\*\*\*\* Nursing Division, Srinagarind Hospital, Faculty of Medicine, Khon Kaen University, Khon Kaen

\*\*\*\*\* Bone and Mineral Research Program, Garvan Institute of Medical Research, Sydney, Australia

\*\*\*\*\* Department of Medicine, Faculty of Medicine, Ramathibodi Hospital, Mahidol University

---

**Background:** Osteoporosis is a growing health problem not only in women but also in men. However, there is a scarcity of epidemiologic data to study osteoporosis in Thai men.

**Objectives:** To examine the bone mineral density (BMD) and to determine the prevalence of osteoporosis in Thai men.

**Material and Method:** A total of 412 men (159 from Bangkok and 253 from Khon Kaen, respectively) averaging  $51 \pm 16$  years of age, were measured for BMD at the femoral neck and lumbar spine by dual energy X-ray absorptiometry (LUNAR Corporation, WI, USA).

**Results:** The peak BMD was observed in men 20-29 years of age at both the femoral neck (mean  $\pm$  SD,  $1.10 \pm 0.15$  g/cm<sup>2</sup>) and lumbar spine (mean  $\pm$  SD,  $1.17 \pm 0.13$  g/cm<sup>2</sup>). The prevalence of osteoporosis in the entire group of subjects was 12.6, 4.6 and 3.9 per cent at the femoral neck, lumbar spine and both sites, respectively. The prevalence of osteoporosis increased with advancing age and was significantly higher at the femoral neck in urban men than rural men (18.2 vs 9.2 per cent,  $p < 0.05$ ) but comparable at the lumbar spine (5.0 vs 4.3 per cent,  $p = 0.81$ ). The correlation between femoral neck and lumbar spine BMDs was 0.53 ( $p < 0.001$ ). In univariate analysis, increased age, lower weight and lesser height were each associated with lower femoral neck BMD, whereas only lower weight and lesser height were associated with lower lumbar spine BMD. However, when the three factors were entered simultaneously, only increased age and lower weight were significantly associated with lower femoral neck BMD and only lower weight had a significant association with lower lumbar spine BMD.

**Conclusion:** The present study demonstrated descriptive BMD data, normal BMD reference values for diagnosis and reported the prevalence of osteoporosis in Thai men.

**Keywords:** Osteoporosis, Bone mineral density, Men, Thai, Epidemiology

**J Med Assoc Thai 2006; 89 (2): 160-9**

**Full text. e-Journal:** <http://www.medassocthai.org/journal>

---

Osteoporosis is a metabolic bone disease characterized by relatively low bone mineral density (BMD), microarchitectural deterioration of bone tissue,

and increased susceptibility to fracture<sup>(1)</sup>. With an aging global population, osteoporosis is fast becoming a worldwide concern because of its age-associated prevalence, costs, morbidity and mortality<sup>(2,3)</sup>. Although, osteoporosis is generally considered a condition affecting postmenopausal women, up to 20 and 30 per cent of symptomatic vertebral and hip fractures, respectively, occur in men<sup>(4-9)</sup>. As the world's popula-

---

Correspondence to : Pongchaiyakul C, Division of Endocrinology and Metabolism, Department of Medicine, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002, Thailand.  
Phone: 0-4336-3664, 0-4336-3746, Fax: 0-4334-7542,  
E-mail: pchatl@kku.ac.th

tion ages, it is predicted that osteoporotic fractures will increase dramatically by the end of this decade<sup>(9)</sup>.

Important sex-specific differences are already known to occur in bone physiology and geometry, fracture epidemiology, bone gonadal hormone response, and post-hip-fracture mortality. These differences point to the importance of doing separate osteoporosis studies in men as it will lead to more specific and effective prevention-based strategies<sup>(10-12)</sup>.

Although several factors contribute to fracture risk, BMD measurement is still the most important element in diagnosing osteoporosis or in screening people at greater risk of fractures<sup>(2)</sup>. The WHO has defined osteoporosis in terms of BMD, based on prior studies using dual energy X-ray absorptiometry (DXA)<sup>(13)</sup>. In order to assure the validity of the results of bone densitometry, those results have to be considered in comparison with the corresponding values that refer to age- and sex-matched, healthy persons from the same population. Although the normal reference data for Thai women has been reported<sup>(14-16)</sup>, there exists no complete set of data regarding the normal range and cut-off values of the hip and spine BMD in Thai men. Therefore, the authors designed a study to examine BMD and to determine the prevalence of osteoporosis in a population-based sample of Thai men.

## **Material and Method**

### ***Setting and Subjects***

The authors designed a cross-sectional study of the Thai population. The Khon Kaen University and Mahidol University Ethics Committees examined and approved the study protocols. Informed written consent was obtained from each participant. The present study was conducted in accordance with the 1975 Helsinki Declaration (revised 1983).

The data was collected in Khon Kaen (North-east-rural) and Bangkok (Central-urban), Thailand. The method of recruitment was previously described in detail<sup>(17)</sup>. Briefly, in Khon Kaen, subjects were recruited from 2 villages in the Muang district. There were 14 hamlets in the two villages. In each hamlet, a full list of subjects was obtained, from which 40 subjects were randomly selected by the village's administrator. The selected subjects were then sent a letter of invitation to participate in the present study. The response rate was 80.3%. In Bangkok, subjects were recruited via a media campaign, and the sampling technique was similar to the scheme used in Khon Kaen, where subjects were randomly selected from 5 districts within the city

of Bangkok. In the present study, the subjects were selected from the databases (159 and 253 from Bangkok and Khon Kaen, respectively), ranging between 20 and 87 years of age.

The authors excluded patients with: bone disorders, chronic diseases or history of taking medications affecting calcium and bone metabolism such as steroids, thyroid hormone, fluoride, bisphosphonates, calcium, antiepileptics, thiazides, calcitonin, alcohol abuse and a previous history of hypogonadism.

### ***Measurements***

Body weight (while wearing light indoor clothing) was measured using an electronic balance (accuracy 0.1 kg) and standing height (without shoes) with a stadiometer (nearest 0.1 cm). Body mass index (BMI) was calculated as the ratio of weight (kg) over height (m<sup>2</sup>).

In Khon Kaen, BMD (g/cm<sup>2</sup>) at the femoral neck and lumbar spine (L2-L4) was measured by DXA using a LUNAR DPX-IQ densitometer (LUNAR Corporation, Madison, WI, USA), while BMDs were measured using a LUNAR DPX-L densitometer (LUNAR Corporation, Madison, WI, USA) in Bangkok. Both study sites used the same protocol. The BMD measurements (from the DXA machines) were corrected using software from the manufacturer.

The coefficient of variation of BMD for normal subjects was 1.5 and 1.3 per cent for the lumbar spine and proximal femur, respectively. The prevalence of osteoporosis was determined by age group. Osteoporosis was defined by a T-score within 2.5 SD or below that of a young adult male designated as the normal reference.

### ***Statistical analyses***

Statistical analyses were performed using SPSS 9.0 (SPSS Inc., Chicago, Illinois). Descriptive results were expressed as the mean, standard deviation (SD) and per cent. Bone mineral density values were analyzed in 10-year intervals by calculating the mean and SD. Normality was confirmed in all age groups. The difference in BMD and prevalence of osteoporosis between urban and rural populations was tested by the unpaired *t*-test and Chi-Square test, respectively. The correlation between BMD, age, body weight and height was obtained using the Pearson correlation coefficient (*r*). Simple linear regression analysis was used to estimate the strength of association between age, weight, height and BMD. The statistical significance was defined at *p* value < 0.05.

## Results

The authors recruited 412 men for the present study. The ratio of subjects from the urban and rural was 2:3. Age averaged  $51 \pm 16$  years (range, 20-87). Other characteristics are shown in Table 1. Body weight was positively associated with BMD at the femoral neck ( $r = 0.14, p = 0.006$ ) and at the lumbar spine ( $r = 0.36, p < 0.001$ ), whereas age was negatively correlated with femoral neck BMD ( $r = 0.51, p < 0.001$ ) but not with lumbar spine BMD ( $r = 0.09, p = 0.07$ ). Height was associated with the femoral neck ( $r = 0.19, p < 0.001$ ) and lumbar spine BMD ( $r = 0.23, p < 0.001$ ). The correlation between femoral neck and lumbar spine BMDs was 0.53 ( $p < 0.001$ ).

The peak BMD was observed in the men between 20 and 29 years of age. Bone mineral density decreased with increasing age; however, the decrease was more pronounced at the femoral neck (Table 2; Fig. 1, 2).

Based on the peak BMD for the entire population, as derived from young adults (mean  $\pm$  SD:  $1.10 \pm 0.15$  for femoral neck and  $1.17 \pm 0.13$  for lumbar spine), the prevalence of osteoporosis was between 12.6 and 4.6 per cent, at the femoral neck and lumbar spine, respectively. Furthermore, it also increased with advancing age: in individuals over 50, 60 and 70, as the prevalence of osteoporosis at the femoral neck was 19.1 (44/230), 23.8 (35/147) and 32.3 (20/62) per cent, respectively, while at the lumbar spine was 7.4 (17/230), 10.2 (14/147) and 14.5 (4/62) per cent, respectively (Table 3).

The age-specific prevalence of osteoporosis between urban and rural subjects was compared and was significantly higher at the femoral neck in urban men ( $p < 0.05$ ), whereas it was not significantly different at the lumbar spine between regions (Table 3).

In the unadjusted analysis, age, weight and height were each associated with femoral neck BMD, whereas only weight and height were associated with lumbar spine BMD. When the three factors were entered simultaneously, only age and weight were significantly associated with femoral neck BMD and only weight had any significant association with lumbar spine BMD. Each one-year increase in age was associated with a  $5.31 \text{ mg/cm}^2$  decrease in femoral neck BMD ( $p < 0.001$ ); and a  $0.79 \text{ mg/cm}^2$  decrease in lumbar spine BMD ( $p = 0.11$ ). Furthermore, each 1-kg increase in weight was associated with a 2.20 increase in the femoral neck BMD ( $p = 0.007$ ) and  $5.48 \text{ mg/cm}^2$  increase in lumbar spine BMD ( $p < 0.001$ ) (Table 4). However, the three factors collectively accounted for 28 and 14 per cent of the variation in femoral neck and lumbar spine BMD, respectively.

## Discussion

Osteoporosis in women has emerged as one of the most common diseases of the elderly and one of the most significant public health issues in the world. This emergence is due in part to the age-related prevalence, and the public appreciation of the seriousness of the consequences vis-à-vis morbidity, economic costs and mortality<sup>(2,18,19)</sup>.

Although less common, osteoporosis in men is also prevalent worldwide with equally serious implications. Specific definitions for male osteoporosis are needed and cost-effective guidelines on who should be investigated and treated and how. The role of BMD measurement in diagnosis and treatment decisions needs to be clarified.

This is the first study done in Thai men recruited from both urban and rural regions. They were examined for BMD and the prevalence of osteoporosis.

**Table 1.** Subject characteristics

	Urban	Rural	Total
Number of subjects	159	253	412
Age (years)	$50.0 \pm 17.5$	$51.4 \pm 16.0$	$50.7 \pm 16.6$
Body weight (kg)	$64.2 \pm 11.1$	$60.8 \pm 10.6$	$62.1 \pm 10.9$
Height (cm)	$165.5 \pm 6.3$	$162.3 \pm 6.3$	$163.6 \pm 6.5$
Body mass index (kg/m <sup>2</sup> )	$23.4 \pm 3.6$	$23.0 \pm 3.1$	$23.1 \pm 3.3$
Bone mineral density (g/cm <sup>2</sup> )			
Femoral neck	$0.87 \pm 0.16$	$0.94 \pm 0.17$	$0.91 \pm 0.17$
Lumbar spine	$1.12 \pm 0.17$	$1.12 \pm 0.17$	$1.12 \pm 0.17$

All values are means  $\pm$  SDs

**Table 2.** Bone mineral density in Thai men by age group

Age group	Femoral neck BMD (g/cm <sup>2</sup> )			Lumbar spine BMD (g/cm <sup>2</sup> )		
	Urban (n = 159)	Rural (n = 253)	Total (n = 412)	Urban (n = 159)	Rural (n = 253)	Total (n = 412)
20-29	1.01 (n = 12)	1.17 (n = 32)	1.10 (n = 57)	1.13 (n = 25)	1.21 (n = 32)	1.17 (n = 57)
30-39	0.94 (n = 28)	1.01 (n = 35)	0.98 (n = 63)	1.16 (n = 28)	1.11 (n = 35)	1.13 (n = 63)
40-49	0.83 (n = 26)	0.96 (n = 36)	0.91 (n = 62)	1.07 (n = 26)	1.07 (n = 36)	1.07 (n = 62)
50-59	0.81 (n = 26)	0.93 (n = 57)	0.89 (n = 83)	1.08 (n = 26)	1.12 (n = 57)	1.11 (n = 83)
60-69	0.83 (n = 28)	0.85 (n = 57)	0.85 (n = 85)	1.12 (n = 28)	1.12 (n = 57)	1.12 (n = 85)
70+	0.80 (n = 26)	0.81 (n = 36)	0.81 (n = 62)	1.14 (n = 26)	1.09 (n = 36)	1.11 (n = 62)

All values are means SDs.


\*Statistical significance at  $p < 0.001$  (Using unpaired *t*-test)

**Table 3.** Prevalence of osteoporosis in urban vs. rural Thai men by age group


Age group	FN			LS			FN or LS			FN and LS		
	Urban	Rural	Total	Urban	Rural	Total	Urban	Rural	Total	Urban	Rural	Total
20-29	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
30-39	-	-	-	-	-	-	-	-	-	-	-	-
40-49	1 (3.6)	-	1 (1.6)	-	1 (2.9)	1 (1.6)	1 (3.6)	1 (2.9)	2 (3.2)	-	-	-
50-59	7 (26.9)**	-	7 (11.3)	-	1 (2.8)	1 (1.6)	7 (26.9)*	1 (2.8)	8 (12.9)	-	-	-
60-69	5 (19.2)	4 (7.0)	9 (10.8)	1 (3.8)	1 (1.8)	2 (2.4)	4 (15.4)	3 (5.3)	7 (8.4)	1 (3.8)	1 (1.8)	2 (2.4)
70+	7 (25.0)	8 (14.0)	15 (17.6)	3 (10.7)	3 (5.3)	6 (7.1)	4 (14.3)	5 (8.8)	9 (10.6)	3 (10.7)	3 (5.3)	6 (7.1)
Total	29 (18.2)*	23 (9.1)	52 (12.6)	8 (5.0)	11 (4.3)	19 (4.6)	21 (13.2)*	18 (7.1)	39 (9.5)	8 (5.0)	8 (3.2)	16 (3.9)

FN; femoral neck, LS; lumbar spine

Statistical significant at  $p < 0.05^*$  and  $< 0.001^{**}$  (Using Chi-Square test)



**Fig. 1** Scatter plot between age and bone mineral density



**Fig. 2** Bone mineral density in urban and rural Thai men by age group

**Table 4.** Regression analyses

	Femoral neck BMD (mg/cm <sup>2</sup> )			Lumbar spine BMD (mg/cm <sup>2</sup> )		
	Coefficients $\pm$ SE <sup>a</sup>	<i>p</i>	<i>R</i> <sup>2.b</sup>	Coefficients $\pm$ SE <sup>a</sup>	<i>p</i>	<i>R</i> <sup>2.b</sup>
Univariate						
Age (per 1 yr)	-5.25 $\pm$ 0.43	<0.001	0.26	-0.90 $\pm$ 0.50	0.072	0.01
Weight (per 1 kg)	2.10 $\pm$ 0.76	0.006	0.02	5.54 $\pm$ 0.71	<0.001	0.13
Height (per 1 cm)	4.95 $\pm$ 1.27	<0.001	0.04	5.93 $\pm$ 1.25	<0.001	0.05
Multivariate						
Age (per 1 yr)	-5.31 $\pm$ 0.45	<0.001	0.28	-0.79 $\pm$ 0.49	0.110	0.14
Weight (per 1 kg)	2.20 $\pm$ 0.81	0.007		5.48 $\pm$ 0.88	<0.001	
Height (per 1 cm)	-0.96 $\pm$ 1.42	0.501		0.09 $\pm$ 1.54	0.954	

<sup>a</sup>Values are regression coefficients  $\pm$  SE describing the change in BMD (mg/cm<sup>2</sup>) associated with a unit change in the factor

<sup>b</sup>Coefficient of determination: the proportion of variation in BMD is explained by the variation in a factor

Osteoporosis (T-score  $\leq -2.5$  SD) at the femoral neck and lumbar spine was diagnosed when the measured BMD was below 0.80 and 0.91 g/cm<sup>2</sup>, respectively. The prevalence of osteoporosis at the femoral neck, lumbar spine and both sites was 12.6, 4.6, and 3.9 per cent, respectively. The prevalence of osteoporosis increased with advancing age. This finding was different from previous studies in Thai women among whom osteoporosis at the lumbar spine was more common than at the femoral neck<sup>(14-16)</sup>. The prevalence of osteoporosis in the present study was consistent with a Taiwanese study<sup>(20)</sup> but different from studies done in Chinese<sup>(21,22)</sup>, Lebanese<sup>(23)</sup> and Caucasian men<sup>(24-30)</sup>.

Yeh et al reported that the prevalence of osteoporosis in men aged 70 and over was 32.8 and 15.7 per cent at the femoral neck and lumbar spine, respectively<sup>(20)</sup>, while Chan et al reported the prevalence of osteoporosis in men was 2.0 and 3.4 per cent at the femoral neck and lumbar spine, respectively, while the prevalence increased to 5.1 per cent in men over 70<sup>(21)</sup>. Malouf et al reported the prevalence of osteoporosis was 9 per cent in both the femoral neck and lumbar spine in Lebanese men<sup>(24)</sup>. Tenenmouse et al reported the prevalence of osteoporosis in Canadian men 50 and over was 6.6 and 2.9 per cent at the femoral neck and lumbar spine, respectively<sup>(25)</sup>. The study from NHANES III showed the prevalence of low BMD was 33 per cent and osteoporosis in white, Hispanic and African-American men 50 years and over was 7, 3 and 5 per cent, respectively<sup>(26)</sup>, while an Australian study showed the prevalence of osteoporosis higher in elderly white men (11 per cent)<sup>(28)</sup>.

Depending on the skeletal site assessed the prevalence of osteoporosis in men varies between 0 and 36 per cent<sup>(12,31)</sup>. The discordance using femoral neck and lumbar spine BMD has been explained, however, different conclusions have been reached concerning the relative sensitivity of hip and spine BMD in the diagnosis of osteoporosis based on the criteria from the WHO.

In some reports, measurements of the femoral neck and total hip identified fewer osteoporotic patients than spine BMD<sup>(32,33)</sup>. While in others, DXA of the hip was a more sensitive indicator for osteoporosis than DXA of the spine<sup>(34,35)</sup>. In the present study, the prevalence of osteoporosis was nominally higher when determined by DXA at the femoral neck *vs* the lumbar spine. This finding may be explained by the process of osteophytosis, which is a natural aging process and usually more prominent at the lumbar spine than the hip, and plays a role in lowering the sensitivity of

DXA in the elderly.

It has been demonstrated that the presence of spinal osteophytes has a major impact on measured BMD in men<sup>(36)</sup> and is associated with a substantial increase in spinal BMD without necessarily any concomitant decrease in fracture risk<sup>(36-39)</sup>. Previous studies reported that spinal BMD, measured by DXA in subjects with osteophytosis, are 15 to 30 per cent higher than age-and-sex matched controls without osteophytosis<sup>(36,37)</sup>. In the present study, the standard deviation was higher in the aging group compared with the younger group (Table 2; Fig. 2).

The incidence of fractures caused by osteoporosis tends to increase with urbanization and the rate of fractures is higher in urban *vs* rural communities<sup>(40-46)</sup>. While many factors are posited as responsible, BMD is thought as the primary determinant, because it is the most consistent and strongest predictor of fracture risk. Indeed, BMD among rural populations is higher than urban populations<sup>(47-49)</sup>; however, most studies focus on Caucasian populations so there is a paucity of data on Asians.

In the present Thai population-based study, the authors observed that BMD in urban men was consistently lower than their rural confreres. The peak bone mass in rural men was significantly higher than in urban men and this finding was consistent with a previous study<sup>(49)</sup>. Secondly, the BMD in rural men was significantly higher than that in an urban population, at the femoral neck (Table 2; Fig. 2) as in previous studies<sup>(47,48)</sup>. The difference in BMD can perhaps be explained by the difference in physical activity between the two day-to-day environments. Rural populations typically have more strenuous, weight-bearing activity than urban populations<sup>(50,51)</sup>. Northeast Thais are mainly farmers who spend their days in manual rice culture or at some other vigorous manual activity.

As in the present study, many prior studies found age and weight correlated to BMD of the hip and lumbar spine in both men and women, while any association between height and BMD was less pronounced<sup>(52,53)</sup>. However, the strength of the association between age, weight and BMD was not strong, particularly at the lumbar spine.

A number of strengths and limitations attend the present study. 1) The data were obtained from large, well-defined rural and urban areas, which made possible reliable delineation of the rural *vs* urban differences. In fact, the subjects in the present study were from two provinces and the authors did not use cluster random sampling for cross-country representa-



tion: such an approach could be used in a follow-on study to test whether the present study represents all Thai men. Notwithstanding, the subjects in the present study were randomly selected and well characterized, thus it is a first important step toward recognizing the prevalence of osteoporosis in men in Thailand. 2) Since the present study focused on Thais, the body size, lifestyle, cultural background and environmental living conditions will differ from other populations; therefore, care should be taken when extrapolating these results to other populations. 3) The measurement error for BMD might result in misclassifications of osteoporosis. 4) Body weight was measured only once, which may not reflect the subject's long-term weight. These sources of measurement errors might skew the results; however, such errors are common in this type of study.

In conclusion, the age-specific and -adjusted prevalence of osteoporosis among Thai men was 12.6 and 4.6 per cent for femoral neck and lumbar spine, respectively. The data provided by the present study could serve as normal reference values for Thai men and for a public health policy to promote bone health and the prevention of osteoporosis in Thai men.

#### Acknowledgments

The present study was supported by the Thai Research Fund. The authors wish to thank Assistant Professor Kanchana Jansoong for her helpful suggestions and Mr. Bryan Roderick Hamman for assistance with the English-language presentation of the manuscript.

#### References

1. Consensus Development Conference: prophylaxis and treatment of osteoporosis. *Am J Med* 1991; 90: 107-10.
2. Kanis JA, Melton LJ III, Christiansen C, Johnston CC, Khaltaev N. The diagnosis of osteoporosis. *J Bone Miner Res* 1994; 9: 1137-41.
3. Cummings SR, Melton LJ. Epidemiology and outcomes of osteoporotic fractures. *Lancet* 2002; 359: 1761-7.
4. Eastell R, Boyle IT, Compston J, Cooper C, Fogelman I, Francis RM, et al. Management of male osteoporosis: report of the UK Consensus Group. *QJM* 1998; 91: 71-92.
5. Pande I, Francis RM. Osteoporosis in men. *Best Pract Res Clin Rheumatol* 2001; 15: 415-27.
6. Bilezikian JP. Osteoporosis in men. *J Clin Endocrinol Metab* 1999; 84: 3431-4.
7. Seeman E. Osteoporosis in men. *Osteoporos Int* 1999; 9(Suppl 2): S97-110.
8. Orwoll ES. Osteoporosis in men. *Endocrinol Metab Clin North Am* 1998; 27: 349-67.
9. Cooper C, Campion G, Melton LJ III. Hip fractures in the elderly a world-wide projection. *Osteoporos Int* 1992; 2: 285-9.
10. Boling EP. Gender and osteoporosis: similarities and sex-specific differences. *J Gend Specif Med* 2001; 4: 36-43.
11. Duan Y, Seeman E. Bone fragility in Asian and Caucasian men. *Ann Acad Med Singapore* 2002; 31: 54-66.
12. Melton LJ III. The prevalence of osteoporosis: gender and racial comparison. *Calcif Tissue Int* 2001; 69: 179-81.
13. Assessment of fracture risk and its application to screening for postmenopausal osteoporosis. Report of a WHO Study Group. *World Health Organ Tech Rep Ser* 1994; 843: 1-129.
14. Pongchaiyakul C, Rojroongwasinkul N, Chotmongkol R, Kosulwat V, Charoenkiatkul S, Rajatanavin R. Bone mineral density in rural Thai adults living in Khon Kaen province. *J Med Assoc Thai* 2002; 85: 235-44.
15. Limpaphayom KK, Taechakraichana N, Jaisamrarn U, Bunyavejchevin S, Chaikittisilpa S, Poshya-chinda M, et al. Prevalence of osteopenia and osteoporosis in Thai women. *Menopause* 2001; 8: 65-9.
16. Taechakraichana N, Angkawanich P, Panyakhamlerd K, Limpaphayom K. Postmenopausal osteoporosis: what is the real magnitude of the problem in the Thai population? *J Med Assoc Thai* 1998; 81: 397-401.
17. Pongchaiyakul C, Nguyen TV, Kosulwat V, Rojroongwasinkul N, Charoenkiatkul S, Rajatanavin R. Effect of urbanization of bone mineral density: a Thai epidemiological study. *BMC Musculoskelet Disord* 2005; 6: 5.
18. Kanis JA. Diagnosis of osteoporosis and assessment of fracture risk. *Lancet* 2002; 359: 1929-36.
19. Cummings SR, Melton LJ. Epidemiology and outcomes of osteoporotic fractures. *Lancet* 2002; 359: 1761-7.
20. Yeh LR, Chen CK, Lai PH. Normal bone mineral density in anteroposterior, lateral spine and hip of Chinese men in Taiwan: effect of age change, body weight and height. *J Chin Med Assoc* 2004; 67: 287-95.
21. Chan WP, Liu JF, Chi WL. Evaluation of bone

- mineral density of the lumbar spine and proximal femur in population-based routine health examinations of healthy Asians. *Acta Radiol* 2004; 45: 59-64.
22. Yu W, Qin M, Xu L. Bone mineral analysis of 445 normal subjects assessed by dual X-ray absorptiometry. *Chin J Radol* 1998; 30: 625-9.
  23. Yu W, Qin M, Xu L. Bone mineral analysis of proximal femur in 445 normal subjects assessed by dual X-ray absorptiometry. *Chin J Radol* 1998; 32: 23-6.
  24. Malouf G, Salem S, Sandid M, Attallah P, Eid J, Saliba N, et al. Bone mineral density of the Lebanese reference population. *Osteoporos Int* 2000; 11: 756-64.
  25. Tenenhouse A, Jorjoseph L, Kreiger N, Poliquin S, Murray TM, Blondeau L, et al. Estimation of the prevalence of low bone density in Canadian women and men using a population-specific DXA reference standard: the Canadian Multicentre Osteoporosis Study (CaMos). *Osteoporos Int* 2000; 11: 897-904.
  26. Looker AC, Orwoll ES, Johnston CC Jr, Lindsay RL, Wahner HW, Dunn WL, et al. Prevalence of low femoral bone density in older US adults from NHANES III. *J Bone Miner Res* 1997; 12: 1761-8.
  27. Melton LJ III, Atkinson EJ, O'Connor MK, O'Fallon WM, Riggs BL. Bone density and fracture risk in men. *J Bone Miner Res* 1998; 13: 1915-23.
  28. Nguyen TV, Center JR, Eisman JA. Osteoporosis in elderly men and women: effects of dietary calcium, physical activity, and body mass index. *J Bone Miner Res* 2000; 15: 322-32.
  29. Kanis JA, Johnell O, Oden A, Jonsson B, De Laet C, Dawson A. Risk of hip fracture according to the World Health Organization criteria for osteopenia and osteoporosis. *Bone* 2000; 27: 585-90.
  30. Kanis JA, Johnell O, Oden A, Sembo I, Redlund-Johnell I, Dawson A, et al. Long-term risk of osteoporotic fracture in Malmo. *Osteoporos Int* 2000; 11: 669-74.
  31. Melton LJ III, Khosla S, Achenbach SJ, O'Connor MK, O'Fallon WM, Riggs BL. Effects of body size and skeletal site on the estimated prevalence of osteoporosis in women and men. *Osteoporos Int* 2000; 11: 977-83.
  32. O'Gradaigh D, Debiram I, Love S, Richards HK, Compston JE. A prospective study of discordance in diagnosis of osteoporosis using spine and proximal femur bone densitometry. *Osteoporos Int* 2003; 14: 13-8.
  33. Arlot ME, Sornay-Rendu E, Garnero P, Vey-Marty B, Delmas PD. Apparent pre- and postmenopausal bone loss evaluated by DXA at different skeletal sites in women: the OFELY cohort. *J Bone Miner Res* 1997; 12: 683-90.
  34. Liu G, Peacock M, Eilam O, Dorulla G, Braunstein E, Johnston CC. Effect of osteoarthritis in the lumbar spine and hip on bone mineral density and diagnosis of osteoporosis in the elderly men and women. *Osteoporos Int* 1997; 7: 564-9.
  35. Rizzoli R, Slosman D, Bonjour JP. The role of dual energy X-ray absorptiometry of lumbar spine and proximal femur in the diagnosis and follow-up of osteoporosis. *Am J Med* 1995; 98: 33S-36S.
  36. Reid IR, Evans MC, Ames R, Wattie DJ. The influence of osteophytes and aortic calcification on spinal mineral density in postmenopausal women. *J Clin Endocrinol Metab* 1991; 72: 1372-4.
  37. Orwoll ES, Oviatt SK, Mann T. The impact of osteophytic and vascular calcifications on vertebral mineral density measurements in men. *J Clin Endocrinol Metab* 1990; 70: 1202-7.
  38. Laitinen K, Valimaki M, Keto P. Bone mineral density measured by dual-energy X-ray absorptiometry in healthy Finnish women. *Calcif Tissue Int* 1991; 48: 224-31.
  39. Masud T, Langley S, Wiltshire P, Doyle DV, Spector TD. Effect of spinal osteophytosis on bone mineral density measurements in vertebral osteoporosis. *BMJ* 1993; 307: 172-3.
  40. Sanders KM, Nicholson GC, Ugoni AM, Seeman E, Pasco JA, Kotowicz MA. Fracture rates lower in rural than urban communities: the Geelong Osteoporosis Study. *J Epidemiol Community Health* 2002; 56: 466-70.
  41. Kaastad TS, Meyer HE, Falch JA. Incidence of hip fracture in Oslo, Norway: differences within the city. *Bone* 1998; 22: 175-8.
  42. Madhok R, Melton LJ III, Atkinson EJ, O'Fallon WM, Lewallen DG. Urban vs rural increase in hip fracture incidence. Age and sex of 901 cases 1980-89 in Olmsted County, U.S.A. *Acta Orthop Scand* 1993; 64: 543-8.
  43. Falch JA, Kaastad TS, Bohler G, Espeland J, Sundsvold OJ. Secular increase and geographical differences in hip fracture incidence in Norway. *Bone* 1993; 14: 643-5.
  44. Norton R, Yee T, Rodgers A, Gray H, MacMahon S. Regional variation in the incidence of hip fracture in New Zealand. *N Z Med J* 1997; 110: 78-80.
  45. Jonsson B, Gardsell P, Johnell O, Redlund-Johnell



- I, Sernbo I. Differences in fracture pattern between an urban and a rural population: a comparative population-based study in southern Sweden. *Osteoporos Int* 1992; 2: 269-73.
46. Mannius S, Mellstrom D, Oden A, Rundgren A, Zetterberg C. Incidence of hip fracture in western Sweden 1974-1982. Comparison of rural and urban populations. *Acta Orthop Scand* 1987; 58: 38-42.
  47. Gardsell P, Johnell O, Nilsson BE, Sernbo I. Bone mass in an urban and a rural population: a comparative, population-based study in southern Sweden. *J Bone Miner Res* 1991; 6: 67-75.
  48. Filip RS, Zagorski J. Bone mineral density and osteoporosis in rural and urban women. Epidemiological study of the Lublin region (Eastern Poland). *Ann Agric Environ Med* 2001; 8: 221-6.
  49. Sundberg M, Duppe H, Gardsell P, Johnell O, Ornstein E, Sernbo I. Bone mineral density in adolescents. Higher values in a rural area - a population-based study of 246 subjects in southern Sweden. *Acta Orthop Scand* 1997; 68: 456-60.
  50. Jonsson B, Gardsell P, Johnell O, Sernbo I, Gullberg B. Life-style and different fracture prevalence: a cross-sectional comparative population-based study. *Calcif Tissue Int* 1993; 52: 425-33.
  51. Ringsberg KA, Gardsell P, Johnell O, Josefsson PO, Obrant KJ. The impact of long-term moderate physical activity on functional performance, bone mineral density and fracture incidence in elderly women. *Gerontology* 2001; 47: 15-20.
  52. Kao CH, Chen CC, Wang SJ. Normal data for lumbar spine bone mineral density in healthy elderly Chinese: influences of sex, age, obesity and ethnicity. *Nucl Med Commun* 1994; 15: 916-20.
  53. Nishizawa Y, Koyama H, Shoji T, Aratani H, Hagiwara S, Miki T, et al. Obesity as a determinant of regional bone mineral density. *J Nutr Sci Vitaminol* 1991; 37(Suppl): S65-70.

---

## ความชุกของโรคกระดูกพรุนในผู้ชายไทย

ฉัตรเลิศ พงษ์ไชยกุล, เฉลิมชัย อภิญาณรักษ์, ศุภศิลาป สุนทราภา, สุกรี สุนทราภา, ชูวงศ์ พงษ์ไชยกุล, ทวน เหมียน, รัชตะ รัชตะนาวิน

**บทนำ:** ปัจจุบันโรคกระดูกพรุนมิได้พบเพียงเฉพาะในเพศหญิงเท่านั้นแต่ยังสามารถพบในเพศชายได้เช่นกัน อย่างไรก็ตามการศึกษาทางระบาดวิทยาในประเทศไทยเกี่ยวกับโรคกระดูกพรุนในผู้ชายยังมีอยู่อย่างจำกัด

**วัตถุประสงค์:** เพื่อศึกษาความหนาแน่นของกระดูกและความชุกของโรคกระดูกพรุนในผู้ชายไทย

**วัสดุและวิธีการ:** มีผู้เข้าร่วมการศึกษาทั้งสิ้นจำนวน 412 คน (159 คนจากกรุงเทพมหานครและ 253 คนจากจังหวัดขอนแก่น) อายุเฉลี่ย  $51 \pm 16$  ปี ทำการวัดความหนาแน่นของกระดูกที่ตำแหน่งกระดูกสะโพกและกระดูกสันหลังระดับเอวที่ 2-4 ด้วยเครื่องวัดความหนาแน่นของกระดูก (LUNAR Corporation, WI, USA)

**ผลการศึกษา:** พบว่ามวลกระดูกสูงสุดอยู่ระหว่างช่วงอายุ 20-29 ปีทั้งที่ตำแหน่งกระดูกสะโพกและกระดูกสันหลังระดับเอว โดยมีค่าเฉลี่ยและค่าเบี่ยงเบนมาตรฐานที่กระดูกสะโพกและกระดูกสันหลังระดับเอวเท่ากับ  $1.10 \pm 0.15$  และ  $1.17 \pm 0.13$  กรัมต่อตารางเซนติเมตร ตามลำดับ พบความชุกของโรคกระดูกพรุนที่ตำแหน่งกระดูกสะโพกร้อยละ 12.6 ที่กระดูกสันหลังระดับเอวร้อยละ 4.6 และทั้งสองตำแหน่งร้อยละ 3.9 โดยพบ ความชุกของโรคกระดูกพรุนเพิ่มขึ้นตามอายุ นอกจากนั้นพบว่าคนที่อาศัยในกรุงเทพมหานครมีความชุกของโรค กระดูกพรุนที่ตำแหน่งกระดูกสะโพกสูงกว่าคนที่อาศัยในจังหวัดขอนแก่น (ร้อยละ 18.2 เทียบกับ 9.2) โดยพบว่ามีค่าแตกต่างอย่างมีนัยสำคัญทางสถิติ ในขณะที่ความชุกของโรคกระดูกพรุนที่ตำแหน่งกระดูกสันหลังระดับเอวระหว่างประชากรสองกลุ่มไม่มีความแตกต่างกัน (ร้อยละ 5 เทียบกับ 4.3) พบความสัมพันธ์ระหว่างความหนาแน่นของกระดูกสะโพกและกระดูกสันหลังระดับเอวเท่ากับ 0.53 โดยมีนัยสำคัญทางสถิติ จากการวิเคราะห์สมการถดถอยอย่างง่ายพบว่าอายุที่เพิ่มขึ้น น้ำหนักตัวและความสูงที่ลดลงมีความสัมพันธ์อย่างมีนัยสำคัญทางสถิติกับการลดลงของความหนาแน่นของกระดูกสะโพก ในขณะที่พบน้ำหนักตัวและความสูงที่ลดลงเพียงสองปัจจัยที่มีความสัมพันธ์อย่างมีนัยสำคัญทางสถิติกับการลดลงของความหนาแน่นของกระดูกสันหลังระดับเอว เมื่อทำการวิเคราะห์สมการถดถอยเชิงพหุพบว่าอายุที่เพิ่มขึ้นและน้ำหนักตัวที่ลดลงมีความสัมพันธ์อย่างมีนัยสำคัญทางสถิติกับการลดลงของความหนาแน่นของกระดูกสะโพก ในขณะที่พบน้ำหนักตัวที่ลดลงเพียงปัจจัยเดียวที่มีความสัมพันธ์อย่างมีนัยสำคัญทางสถิติกับการลดลงของความหนาแน่นของกระดูกสันหลังระดับเอว

**สรุป:** การศึกษานี้ได้นำเสนอข้อมูลเชิงพรรณนาของความหนาแน่นของกระดูก ได้แสดงค่ามาตรฐานของความหนาแน่นของกระดูกจากประชากรอายุน้อยที่ใช้ในการเปรียบเทียบเพื่อการวินิจฉัยโรคกระดูกพรุนรวมทั้งได้ศึกษาความชุกของโรคกระดูกพรุนในผู้ชายไทย